


Annual Performance Measure Update: 2009-2016



**Sue O'Brien RN, MSN, ANVP
Nurse Consultant
MOSAIC
Heart Disease & Stroke Prevention**

VTE Prophylaxis STK- 1

Percent of patients with an ischemic stroke, or a hemorrhagic stroke, or stroke not otherwise specified who receive VTE prophylaxis the day of or the day after hospital admission.




Antithrombotic at Discharge STK-2

Percent of patients with an ischemic stroke or TIA prescribed antithrombotic therapy at discharge.



National: 98.5 %

MI: 98.8%




Anticoagulation for A-Fib/A-Flutter STK-3

Percent of patients with an ischemic stroke or TIA with atrial fibrillation/flutter discharged on anticoagulation therapy.



National: 96.3%
MI: 93.3%




IV rt-PA : Arrive by 2 hour, Treat by 3 Hour STK-4

Percent of acute ischemic stroke patients who arrive at the hospital within 120 minutes (2 hours) of time last known well and for whom IV t-PA was initiated at this hospital within 180 minutes (3 hours) of time last known well.


National: 88.1 %

MI: 80.7%



Early Antithrombotics STK-5

Percent of patients with ischemic stroke or TIA who receive antithrombotic therapy by the end of hospital day two.




LDL 100 or ND – Statin STK-6

Percent of Ischemic stroke or TIA patients with LDL = 100, or LDL not measured, or on cholesterol-reducer prior to admission who are discharged on Statin Medication.



National: 96.7%

MI: 96.2%




Dysphagia Screen STK-7

Percent of Stroke patients who undergo screening for dysphagia with an evidence-based bedside testing protocol approved by the hospital before being given any food, fluids, or medication by mouth.




National: 84.8%

MI: 76.8%



Stroke Education STK-8

Percent of patients with Stroke or TIA or their caregivers who were given education and/or educational materials during the hospital stay addressing ALL of the following: personal risk factors for stroke, warning signs for stroke, activation of emergency medical system, need for follow-up after discharge, and medications prescribed.




Smoking Cessation STK-9

Percent of patients with ischemic or hemorrhagic stroke, or TIA with a history of smoking cigarettes, who are, or whose caregivers are, given smoking cessation advice or counseling during hospital stay.




National: 97.6%

MI: 94.5%



Assessed for Rehabilitation STK-10

Percent of patients with Stroke who were assessed for rehabilitation services.



Defect Free Care

Defect-Free Measure of the 10 Consensus CDC/COV measures



National: 81.1%

MI: 75.5 %

