

**Estimates for Selected Risk Factors and Health Indicators
by Emergency Preparedness Region**

State of Michigan

**Michigan Behavioral Risk Factor Surveillance System
(Michigan BRFSS)**

2019-2021

Lifecourse Epidemiology and Genomics Division
Bureau of Epidemiology and Population Health
Michigan Department of Health and Human Services

Table of Contents

Introduction	3
Map of Emergency Preparedness Regions	4
Health Status Tables	
Table 1: General Health Status	5
Table 2: Health Status on at Least 14 Days in the Past Month	6
Table 3: Mean Number of Days of Poor Physical Health, Poor Mental Health, and Activity Limitations.....	7
Table 4: Disability	8
Table 5: Weight Status	9
Table 6: Health Care Coverage Among Adults 18-64 Years Old	10
Table 7: Health Care Access	11
Risk Behaviors Tables	
Table 8: No Leisure Time Physical Activity	12
Table 9: Cigarette Smoking	13
Table 10: Current Smokers Who Attempted to Quit	14
Table 11: E-Cig Use	15
Table 12: Hookah Use	16
Table 13: Alcohol Consumption.....	17
Table 14: Mean Number of Binge Drinking Occassions During the Past Month Among Binge Drinkers	18
Table 15: Hypertension Awareness and Medication Use	19
Clinical Preventative Practices	
Table 16: Cholesterol Screening and Awareness	20
Table 17: No Routine Checkup in Past Year.....	21
Table 18: Immunizations Among Adults Aged 65 Years and Older	22
Table 19: HIV Testing Among Adults Aged 18 - 64 Years	23
Chronic Conditions	
Table 20: Asthma Among Adults	24
Table 21: Asthma Among Children.....	25
Table 22: Chronic Obstructive Pulmonary Disease (COPD).....	26
Table 23: Arthritis.....	27
Table 24: Cardiovascular Disease.....	28
Table 25: Cancer	29
Table 26: Diabetes	30
Table 27: Kidney Disease.....	31
Table 28: Depression	32

Introduction

The Michigan Behavioral Risk Factor Survey (MiBRFS) is an annual, statewide telephone survey of Michigan adults aged 18 years and older that is conducted through a collaborative effort among the Population Health Surveillance Branch (PHSB) of the Centers for Disease Control and Prevention (CDC), the Michigan State University Institute for Public Policy and Social Research (IPPSR), and the Michigan Department of Health and Human Services. Michigan BRFSS data contribute to the national [Behavioral Risk Factor Surveillance System](#) (BRFSS) that is managed by the PHSB at the CDC.

For this report, data from the 2019-2021 Michigan BRFSS were combined. The total sample sizes by year were: 10,518 in 2019, 7,269 in 2020, and 9,425 in 2021. Not all questions were included annually in the survey. The table subtitles indicate which years were used for that particular analysis.

For the 2019 through 2021 Michigan Behavioral Risk Factor Surveys, the sample of land line telephone numbers that were utilized for data collection was selected using a list-assisted, random-digit-dialed methodology with a disproportionate stratification based on phone bank density, and whether or not the phone numbers were directory listed. The sample of cell phone numbers used within each survey year was randomly selected from dedicated cellular telephone banks sorted on the basis of area code and exchange within the State of Michigan.

Iterative proportional fitting or raking was the weighting methodology used to make the 2019-2021 Michigan BRFSS data representative of the current Michigan adult population. Estimates based on this methodology were weighted to adjust for the probabilities of selection and a raking adjustment factor that adjusted for the distribution of Michigan adults by telephone source (land line vs. cell phone), detailed race/ethnicity, education level, marital status, age by gender, gender by race/ethnicity, age by race/ethnicity, and renter/owner status. No additional weighting factors were computed at the regional level, thus the weights used might not always provide accurate regional estimates, particularly when regional demographic distributions vary greatly from that of the state.

In 2021, Michigan BRFSS data from 2019-2021 were analyzed by Michigan Emergency Preparedness (EP) region. A map identifying each of Michigan's eight emergency preparedness regions is included on the following page. Population-based prevalence estimates and asymmetric confidence intervals were calculated for indicators of health status, health risk behaviors, clinical preventive practices, and chronic conditions among the adult population in Michigan. All analyses were performed in SAS-Callable SUDAAN®, a statistical computing that is designed for complex sample surveys.

In an effort to provide the most reliable information possible, prevalence estimates are suppressed if they are based on a denominator of less than 50 respondents or have a relative standard error of greater than 30%.

Due to the BRFSS methodology changes that took place in 2011, the estimates provided within the following tables should not be compared to Michigan BRFSS estimates from years prior to 2011. Methodology changes also took place in 2017 for Fruit and Vegetable Consumption; therefore, these variables were excluded from multi-year analysis.

If you have any questions regarding any of the estimates within the following tables, please contact Yan Tian at TianY@michigan.gov.

Local Health Jurisdictions, Emergency Preparedness Regions and Michigan Counties




Table 1: Health Status by EP Region Michigan BRFSS, 2019-2021 Combined		
	General Health, Fair or Poor^a	
	%	95% Confidence Interval
Michigan Total	16.9	(16.3-17.5)
EP Region		
Region 1	15.4	(14.0-17.0)
Region 2N	14.5	(13.3-15.9)
Region 2S	18.7	(17.3-20.1)
Region 3	20.4	(18.5-22.5)
Region 5	18.8	(16.8-21.1)
Region 6	15.4	(13.8-17.2)
Region 7	16.3	(13.8-19.3)
Region 8	17.8	(14.9-21.3)
^a Among all adults, the proportion who reported that their health, in general, was either fair or poor. (N = 27,155)		

Table 2: Health Status on at Least 14 Days in the Past Month by EP Region Michigan BRFSS, 2019-2021 Combined						
	Poor Physical Health^a		Poor Mental Health^b		Activity Limitation^c	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	12.6	(12.1-13.2)	15.2	(14.6-15.8)	9.4	(9.0-9.9)
EP Region						
Region 1	12.5	(11.2-14.0)	15.5	(14.0-17.2)	8.9	(7.7-10.1)
Region 2N	10.8	(9.8-11.9)	14.2	(12.9-15.5)	8.2	(7.3-9.3)
Region 2S	13.3	(12.1-14.6)	16.4	(15.1-17.8)	10.4	(9.4-11.5)
Region 3	15.5	(13.8-17.3)	16.5	(14.7-18.4)	11.6	(10.1-13.2)
Region 5	14.0	(12.2-16.0)	16.5	(14.5-18.8)	10.3	(8.7-12.3)
Region 6	12.0	(10.6-13.7)	13.5	(12.1-15.1)	8.5	(7.4-9.8)
Region 7	10.9	(8.9-13.2)	14.0	(11.6-16.8)	9.5	(7.5-12.0)
Region 8	14.2	(11.3-17.6)	12.2	(9.6-15.4)	9.3	(7.2-12.0)

^a Among all adults, the proportion who reported 14 or more days of poor physical health, which includes physical illness and injury, during the past 30 days. (N = 26,690)

^b Among all adults, the proportion who reported 14 or more days of poor mental health, which includes stress, depression, and problems with emotions, during the past 30 days. (N = 26,758)

^c Among all adults, the proportion who reported 14 or more days in the past 30 days in which either poor physical health or poor mental health kept respondents from doing their usual activities, such as self-care, work, and recreation. (N = 26,925)

Table 3: Mean Number of Days of Poor Physical Health, Poor Mental Health, and Activity Limitations by EP Region
Michigan BRFSS, 2019-2021 Combined

	Poor Physical Health ^a		Poor Mental Health ^b		Activity Limitation ^c	
	Mean # of days	95% Confidence Interval	Mean # of days	95% Confidence Interval	Mean # of days	95% Confidence Interval
Michigan Total	3.9	(3.8-4.1)	5.0	(4.9-5.1)	3.0	(2.9-3.1)
EP Region						
Region 1	3.9	(3.5-4.2)	5.1	(4.7-5.5)	2.8	(2.5-3.1)
Region 2N	3.4	(3.1-3.7)	4.7	(4.4-5.0)	2.8	(2.6-3.1)
Region 2S	4.1	(3.8-4.4)	5.3	(5.0-5.6)	3.2	(3.0-3.5)
Region 3	4.6	(4.2-5.0)	5.5	(5.0-6.0)	3.6	(3.2-3.9)
Region 5	4.3	(3.8-4.8)	5.4	(4.8-5.9)	3.1	(2.7-3.5)
Region 6	3.8	(3.4-4.2)	4.6	(4.2-5.0)	2.7	(2.5-3.0)
Region 7	4.0	(3.4-4.6)	4.6	(4.0-5.3)	2.8	(2.2-3.3)
Region 8	4.5	(3.7-5.3)	4.4	(3.6-5.1)	2.8	(2.3-3.4)

^a Among all adults, the mean number of days during the past 30 days in which physical health, including illness and injury, was not good. (N = 26,690)

^b Among all adults, the mean number of days during the past 30 days in which mental health, including stress, depression, and problems with emotions, was not good. (N = 26,758)

^c Among all adults, the mean number of days during the past 30 days in which poor physical health or poor mental health kept respondent from doing their usual activities, such as self-care, work, and recreation. (N = 26,925)

Table 4: Disability by EP Region Michigan BRFSS, 2019-2021 Combined		
	Total Disability^a	
	%	95% Confidence Interval
Michigan Total	28.9	(28.1-29.6)
EP Region		
Region 1	28.7	(26.8-30.7)
Region 2N	24.5	(23.0-26.1)
Region 2S	30.5	(28.8-32.2)
Region 3	34.8	(32.5-37.2)
Region 5	31.5	(29.0-34.0)
Region 6	27.4	(25.4-29.4)
Region 7	32.9	(29.3-36.7)
Region 8	29.5	(25.6-33.7)
^a Among all adults, the proportion who reported having serious difficulty hearing, visual impairment, difficulty concentrating, remembering, or making decisions, difficulty walking or climbing stairs, difficulty dressing or bathing, or difficulty doing errands alone. (N = 26,086)		

Table 5: Weight Status by EP Region
Michigan BRFSS, 2019-2021 Combined

	Obese ^a		Overweight ^b		Healthy Weight ^c		Underweight ^d	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	35.2	(34.4-36.0)	34.4	(33.6-35.2)	28.5	(27.8-29.3)	1.9	(1.6-2.2)
EP Region								
Region 1	35.0	(32.9-37.1)	34.2	(32.1-36.3)	29.0	(27.0-31.1)	1.8	(1.3-2.7)
Region 2N	32.2	(30.5-33.9)	35.9	(34.1-37.7)	29.9	(28.2-31.6)	2.1	(1.5-2.9)
Region 2S	36.4	(34.7-38.2)	33.3	(31.6-35.1)	28.4	(26.8-30.1)	1.8	(1.4-2.5)
Region 3	40.6	(38.1-43.2)	33.7	(31.4-36.1)	23.0	(20.9-25.3)	2.7	(1.8-4.0)
Region 5	37.6	(35.0-40.3)	33.4	(31.0-36.0)	27.7	(25.3-30.2)	1.2	(0.8-2.0)
Region 6	34.9	(32.9-37.1)	35.5	(33.4-37.7)	28.3	(26.3-30.5)	1.2	(0.8-1.7)
Region 7	33.3	(29.9-36.9)	33.9	(30.6-37.5)	30.5	(27.1-34.1)	2.3	(1.3-3.8)
Region 8	35.0	(30.8-39.4)	34.5	(30.2-39.0)	30.1	(26.1-34.4)	-- ^e	-- ^e

Note: BMI, body mass index, is defined as weight (in kilograms) divided by height (in meters) squared [weight in kg/(height in meters)²]. Weight and height were self-reported. Pregnant women were excluded. (N = 24,934)

^a Among all adults, the proportion of respondents whose BMI was greater than or equal to 30.0.

^b Among all adults, the proportion of respondents whose BMI was greater than or equal to 25.0, but less than 30.0.

^c Among all adults, the proportion of respondents whose BMI was greater than or equal to 18.5, but less than 25.0.

^d Among all adults, the proportion of respondents whose BMI was less than 18.5.

^e Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 6: Health Care Coverage Among Adults 18-64 Years Old by EP Region Michigan BRFSS, 2019-2021 Combined		
	No Health Care Coverage^a (Among 18 - 64 year olds)	
	%	95% Confidence Interval
Michigan Total	8.6	(8.1-9.2)
EP Region		
Region 1	7.8	(6.5-9.4)
Region 2N	6.3	(5.3-7.5)
Region 2S	8.7	(7.4-10.0)
Region 3	9.1	(7.5-11.0)
Region 5	7.8	(6.2-9.8)
Region 6	9.4	(8.0-11.1)
Region 7	13.0	(8.9-18.6)
Region 8	10.1	(7.2-13.8)
^a Among adults aged 18-64 years, the proportion who reported having no health care coverage, including health insurance, prepaid plans such as HMOs, or government plans, such as Medicare or Indian Health Services. (N = 16,506)		

Table 7: Health Care Access by EP Region Michigan BRFSS, 2019-2021 Combined				
	No Personal Health Care Provider^a		No Health Care Access During Past 12 Months Due to Cost^b	
	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	13.3	(12.7-13.9)	9.2	(8.7-9.7)
EP Region				
Region 1	14.6	(13.0-16.3)	8.8	(7.6-10.1)
Region 2N	10.2	(9.1-11.5)	8.4	(7.4-9.5)
Region 2S	14.4	(13.0-15.8)	9.6	(8.6-10.7)
Region 3	14.1	(12.3-16.1)	10.8	(9.3-12.5)
Region 5	12.6	(10.8-14.6)	8.6	(7.1-10.3)
Region 6	11.8	(10.3-13.5)	9.7	(8.4-11.2)
Region 7	13.3	(10.3-17.1)	7.2	(5.5-9.3)
Region 8	11.7	(9.2-14.7)	7.4	(5.5-9.9)
^a Among all adults, the proportion who reported that they did not have anyone that they thought of as their personal doctor or health care provider. (N = 27,000)				
^b Among all adults, the proportion who reported that in the past 12 months, they could not see a doctor when they needed to due to the cost. (N = 27,141)				

Table 8: No Leisure Time Physical Activity by EP Region Michigan BRFSS, 2019-2021 Combined		
	No Leisure Time Physical Activity^a	
	%	95% Confidence Interval
Michigan Total	23.1	(22.4-23.8)
EP Region		
Region 1	22.7	(20.9-24.5)
Region 2N	20.0	(18.6-21.5)
Region 2S	25.1	(23.5-26.7)
Region 3	27.7	(25.6-30.0)
Region 5	24.4	(22.2-26.7)
Region 6	20.5	(18.9-22.3)
Region 7	25.5	(21.9-29.3)
Region 8	24.1	(20.7-27.9)

^a Among all adults, the proportion who reported not participating in any leisure time physical activities or exercises such as running, calisthenics, golf, gardening, or walking during the past month. (N = 26,856)

Table 9: Cigarette Smoking by EP Region
Michigan BRFSS, 2019-2021 Combined

	Current Smoking ^a		Former Smoking ^b		Never Smoked	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	18.0	(17.4-18.7)	26.5	(25.8-27.2)	55.5	(54.7-56.3)
EP Region						
Region 1	17.1	(15.5-18.9)	27.7	(25.9-29.6)	55.2	(53.0-57.3)
Region 2N	14.7	(13.4-16.1)	27.3	(25.8-28.9)	58.0	(56.2-59.8)
Region 2S	18.8	(17.4-20.3)	23.4	(22.0-24.9)	57.8	(56.0-59.6)
Region 3	22.8	(20.7-25.0)	29.1	(26.9-31.4)	48.2	(45.6-50.7)
Region 5	20.3	(18.0-22.8)	27.6	(25.4-29.9)	52.1	(49.4-54.7)
Region 6	18.0	(16.1-20.0)	25.5	(23.7-27.3)	56.5	(54.3-58.7)
Region 7	18.8	(15.8-22.2)	31.1	(27.8-34.6)	50.1	(46.2-53.9)
Region 8	20.6	(17.1-24.5)	28.8	(25.2-32.8)	50.6	(46.3-54.8)

^a Among all adults, the proportion who reported that they had ever smoked at least 100 cigarettes (5 packs) in their life and that they smoke cigarettes now, either every day or on some days. (N = 26,040)

^b Among all adults, the proportion who reported that they had ever smoked at least 100 cigarettes (5 packs) in their life, but they do not smoke cigarettes now. (N = 26,040)

Table 10: Current Smokers Who Attempted to Quit by EP Region Michigan BRFSS, 2019-2021 Combined		
	Tried to Quit Smoking One Day or Longer in Past Year^a	
	%	95% Confidence Interval
Michigan Total	57.5	(55.3-59.7)
EP Region		
Region 1	55.0	(49.0-60.8)
Region 2N	59.0	(53.5-64.2)
Region 2S	62.3	(57.5-66.9)
Region 3	59.2	(53.5-64.7)
Region 5	57.1	(49.9-64.0)
Region 6	55.3	(48.7-61.7)
Region 7	48.4	(38.8-58.0)
Region 8	53.9	(43.0-64.4)

^a Among current smokers, the proportion who reported that during the past 12 months, they had tried to quit smoking for one day or longer. (N = 3,355)

Table 11: E-Cigarette Use Michigan BRFSS, 2019-2021 Combined		
	Current E-Cigarette Use^a	
	%	95% Confidence Interval
Michigan Total	6.7	(6.0-7.5)
EP Region		
Region 1	7.1	(5.4-9.2)
Region 2N	8.1	(6.5-10.0)
Region 2S	6.4	(4.9-8.2)
Region 3	6.9	(5.1-9.2)
Region 5	10.0	(7.3-13.5)
Region 6	3.8	(2.6-5.6)
Region 7	3.4	(2.0-5.9)
Region 8	-- ^b	-- ^b
^a Among all adults, the proportion who reported that they currently use e-cigarettes or other electronic vaping products, either every day or on some days. (N = 10,515) ^b Suppressed due to a denominator < 50 and/or a relative standard error > 30%.		

Table 12: Current Hookah Use by EP Region Michigan BRFSS, 2019-2021 Combined		
	Current Hookah Use^a	
	%	95% Confidence Interval
Michigan Total	2.7	(2.3-3.1)
EP Region		
Region 1	1.8	(1.2-2.9)
Region 2N	2.7	(1.9-3.7)
Region 2S	3.8	(3.0-5.0)
Region 3	2.1	(1.4-3.2)
Region 5	2.8	(1.7-4.5)
Region 6	2.1	(1.5-3.1)
Region 7	-- ^b	-- ^b
Region 8	-- ^b	-- ^b

^a Among all adults, the proportion who reported smoking tobacco using a hookah, narghile, or water pipe on one or more days during the previous thirty day. (N = 15,562)

^b Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 13: Alcohol Consumption by EP Region
Michigan BRFSS, 2019-2021 Combined

	Any Alcohol in Past Month ^a		Heavy Drinking ^b		Binge Drinking ^c	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	55.3	(54.5-56.1)	6.5	(6.1-6.9)	17.2	(16.6-17.8)
EP Region						
Region 1	55.8	(53.6-57.9)	5.9	(5.0-7.0)	18.4	(16.6-20.2)
Region 2N	60.3	(58.4-62.0)	6.1	(5.3-7.0)	17.1	(15.8-18.6)
Region 2S	53.1	(51.3-54.9)	5.9	(5.1-6.8)	16.7	(15.3-18.1)
Region 3	51.7	(49.2-54.2)	5.9	(4.8-7.2)	17.3	(15.3-19.6)
Region 5	52.9	(50.3-55.6)	6.5	(5.3-7.9)	14.2	(12.4-16.2)
Region 6	55.5	(53.2-57.7)	7.7	(6.5-9.1)	17.6	(15.9-19.4)
Region 7	54.3	(50.4-58.2)	7.5	(5.9-9.5)	15.2	(12.8-17.9)
Region 8	57.4	(53.1-61.6)	9.7	(7.3-12.9)	22.3	(18.7-26.3)

^a Among all adults, the proportion who reported consuming at least one drink of any alcoholic beverage in the previous month. (N = 25,881)

^b Among all adults, the proportion who reported consuming an average of more than two alcoholic drinks per day for men or more than one per day for women in the previous month. (N = 25,657)

^c Among all adults, the proportion who reported consuming five or more drinks per occasion (for men) or 4 or more drinks per occasion (for women) at least once in the previous month. (N = 25,674)

Table 14: Mean Number of Binge Drinking Occasions during the Past Month among Binge Drinkers by EP Region Michigan BRFSS, 2019-2021 Combined		
	Number of Binge Drinking Occasions^a	
	Mean	95% Confidence Interval
Michigan Total	4.9	(4.6-5.2)
EP Region		
Region 1	4.2	(3.6-4.7)
Region 2N	4.2	(3.6-4.8)
Region 2S	4.8	(4.1-5.5)
Region 3	4.6	(3.8-5.3)
Region 5	5.1	(4.1-6.1)
Region 6	5.9	(4.9-7.0)
Region 7	6.7	(4.9-8.6)
Region 8	5.8	(4.4-7.3)

^a Among adults who binge drank in the past month, the mean number of times during which five or more alcoholic beverages (for men) or four or more (for women) were consumed during the past month. (N = 3,650)

	Ever Told HBP ^a		Taking BP Medication ^b	
	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	35.2	(34.4-36.1)	77.9	(76.6-79.2)
EP Region				
Region 1	34.7	(32.4-37.1)	76.4	(72.4-79.9)
Region 2N	33.6	(31.8-35.5)	81.0	(78.2-83.5)
Region 2S	36.3	(34.4-38.2)	78.1	(75.0-80.9)
Region 3	40.8	(38.2-43.5)	82.4	(78.8-85.6)
Region 5	38.4	(35.5-41.3)	78.4	(73.6-82.5)
Region 6	32.9	(30.7-35.1)	72.8	(68.4-76.7)
Region 7	36.3	(32.6-40.1)	77.9	(72.6-82.5)
Region 8	36.3	(31.7-41.1)	79.8	(72.8-85.3)

^a Among all adults, the proportion who reported that they were ever told by a doctor that they had high blood pressure (HBP). Women who had HBP only during pregnancy and adults who were borderline hypertensive were considered to not have been diagnosed. (N = 19,863)

^b Among adults who were ever told that they had HBP, the proportion that reported they were currently taking blood pressure (BP) medicines for their HBP. (N = 8,038)

Table 16: Cholesterol Screening and Awareness by EP Region
Michigan BRFSS, 2019 and 2021 Combined

	Cholesterol Ever Checked ^a		Checked Within Past Five Years ^b		Ever Told High Cholesterol ^c	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	92.6	(92.0-93.1)	89.1	(88.4-89.7)	34.5	(33.6-35.3)
EP Region						
Region 1	91.0	(89.0-92.7)	87.8	(85.7-89.7)	34.0	(31.7-36.4)
Region 2N	95.2	(94.0-96.1)	92.9	(91.6-94.0)	35.8	(33.9-37.8)
Region 2S	93.7	(92.4-94.7)	90.5	(89.1-91.8)	32.6	(30.7-34.5)
Region 3	92.4	(90.5-93.9)	88.3	(86.0-90.2)	37.2	(34.6-39.9)
Region 5	90.4	(87.9-92.5)	87.5	(84.9-89.7)	35.7	(32.9-38.7)
Region 6	92.8	(91.0-94.2)	88.2	(86.0-90.0)	32.6	(30.3-34.9)
Region 7	92.5	(89.7-94.5)	88.5	(85.2-91.1)	39.9	(35.8-44.1)
Region 8	88.5	(83.4-92.2)	81.8	(76.6-86.1)	37.5	(32.6-42.7)

^a Among all adults, the proportion who reported ever having had their blood cholesterol checked. (N = 18,594)

^b Among all adults, the proportion who reported that they have had their blood cholesterol checked within the past five years. (N = 18,594)

^c Among adults who ever had their blood cholesterol checked, the proportion who reported that a doctor, nurse, or other health professional had told them that their cholesterol was high. (N = 18,208)

Table 17: No Routine Checkup in Past Year by EP Region Michigan BRFSS, 2019-2021 Combined		
	Had No Routine Checkup in Past Year^a	
	%	95% Confidence Interval
Michigan Total	22.0	(21.3-22.7)
EP Region		
Region 1	23.2	(21.4-25.2)
Region 2N	20.6	(19.1-22.1)
Region 2S	20.9	(19.5-22.4)
Region 3	21.5	(19.5-23.7)
Region 5	23.2	(21.0-25.6)
Region 6	23.8	(21.9-25.9)
Region 7	20.4	(17.0-24.3)
Region 8	22.1	(18.7-26.0)

^a Among all adults, the proportion who reported that they did not have a routine checkup in the past year. (N = 26,913)

**Table 18: Immunizations among Adults Aged 65 Years and Older
by EP Region
Michigan BRFSS, 2019-2021 Combined**

	Had Flu Vaccine in Past Year ^a		Ever Had Pneumonia Vaccine ^b	
	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	67.5	(66.2-68.8)	72.1	(70.8-73.4)
EP Region				
Region 1	70.5	(67.3-73.5)	77.4	(74.4-80.1)
Region 2N	71.5	(68.8-74.2)	74.6	(71.9-77.2)
Region 2S	64.1	(60.9-67.2)	67.5	(64.2-70.7)
Region 3	61.7	(57.7-65.6)	68.4	(64.3-72.1)
Region 5	70.8	(67.0-74.4)	73.4	(69.5-77.0)
Region 6	69.3	(66.0-72.4)	74.9	(71.6-77.8)
Region 7	68.5	(62.3-74.2)	70.0	(63.9-75.5)
Region 8	63.1	(56.3-69.5)	73.1	(66.0-79.2)

^a Among adults aged 65 years and older, the proportion who reported that they had a flu vaccine, either by an injection in the arm or sprayed in the nose during the past 12 months. (N = 9,348)

^b Among adults aged 65 years and older, the proportion who reported that they ever had a pneumococcal vaccine. (N = 9,058)

Table 19: HIV Testing among Adults Aged 18 - 64 Years by EP Region Michigan BRFSS, 2019-2021 Combined		
	Ever Had an HIV Test^a	
	%	95% Confidence Interval
Michigan Total	44.9	(43.8-45.9)
EP Region		
Region 1	44.2	(41.4-46.9)
Region 2N	43.3	(41.0-45.6)
Region 2S	52.5	(50.3-54.7)
Region 3	41.8	(38.6-45.0)
Region 5	47.5	(44.0-50.9)
Region 6	41.2	(38.4-44.0)
Region 7	38.0	(33.3-42.9)
Region 8	40.3	(35.0-45.9)

^a Among adults aged 18 - 64 years, the proportion who reported that they ever had been tested for HIV, apart from tests that were part of a blood donation. Don't know was considered a valid response to this question. (N = 14,890)

Table 20: Asthma among Adults by EP Region Michigan BRFSS, 2019-2021 Combined				
	Lifetime Asthma Prevalence^a		Current Asthma Prevalence^b	
	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	16.0	(15.4-16.6)	11.2	(10.7-11.8)
EP Region				
Region 1	16.8	(15.2-18.5)	11.9	(10.6-13.4)
Region 2N	17.0	(15.6-18.4)	11.3	(10.2-12.5)
Region 2S	16.6	(15.4-18.0)	12.4	(11.3-13.5)
Region 3	16.0	(14.2-17.9)	11.8	(10.2-13.6)
Region 5	16.7	(14.8-18.9)	11.8	(10.1-13.6)
Region 6	14.5	(13.0-16.2)	9.7	(8.5-11.1)
Region 7	16.6	(13.4-20.4)	11.9	(9.0-15.6)
Region 8	14.0	(11.4-17.1)	9.1	(7.1-11.5)
^a Among all adults, the proportion who reported that they were ever told by a doctor, nurse, or other health care professional that they had asthma. (N = 27,113) ^b Among all adults, the proportion who reported that they still had asthma. (N = 26,968)				

Table 21: Asthma among Children by EP Region Michigan BRFSS, 2019-2021 Combined				
	Lifetime Asthma Prevalence^a		Current Asthma Prevalence^b	
	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	11.2	(10.2-12.4)	7.7	(6.8-8.8)
EP Region				
Region 1	13.1	(10.2-16.5)	8.7	(6.4-11.7)
Region 2N	11.3	(9.1-14.0)	7.0	(5.2-9.3)
Region 2S	12.8	(10.4-15.6)	8.8	(6.9-11.1)
Region 3	12.7	(9.7-16.6)	9.2	(6.5-12.9)
Region 5	9.8	(6.9-13.7)	7.6	(5.0-11.3)
Region 6	9.4	(7.2-12.3)	6.6	(4.7-9.3)
Region 7	-- ^c	-- ^c	-- ^c	-- ^c
Region 8	-- ^c	-- ^c	-- ^c	-- ^c
^a Estimated proportion of Michigan children aged 0-17 years who were ever told by a doctor, nurse or other health professional that they had asthma, using proxy information from the adult respondent. (N = 5,180) ^b Estimated proportion of Michigan children aged 0-17 years who still have asthma, using proxy information from the adult respondent. (N = 5,157) ^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.				

Table 22: Chronic Obstructive Pulmonary Disease (COPD) by EP Region Michigan BRFSS, 2019-2021 Combined		
	Ever Told COPD, Emphysema or Chronic Bronchitis^a	
	%	95% Confidence Interval
Michigan Total	8.1	(7.7-8.6)
EP Region		
Region 1	7.9	(6.9-9.1)
Region 2N	7.3	(6.4-8.2)
Region 2S	8.5	(7.6-9.5)
Region 3	11.3	(9.9-13.0)
Region 5	8.2	(6.8-9.8)
Region 6	6.9	(6.1-7.9)
Region 7	8.5	(6.7-10.6)
Region 8	8.6	(6.5-11.3)

^a Among all adults, the proportion who reported ever being told by a doctor that they had chronic obstructive pulmonary disease (COPD), emphysema or chronic bronchitis. (N = 27,078)

Table 23: Arthritis by EP Region Michigan BRFSS, 2019-2021 Combined		
	Ever Told Arthritis^a	
	%	95% Confidence Interval
Michigan Total	30.3	(29.6-30.9)
EP Region		
Region 1	32.0	(30.2-34.0)
Region 2N	28.7	(27.2-30.2)
Region 2S	29.8	(28.3-31.4)
Region 3	36.4	(34.2-38.7)
Region 5	32.0	(29.7-34.4)
Region 6	28.5	(26.7-30.3)
Region 7	32.6	(29.3-36.0)
Region 8	36.3	(32.4-40.4)

^a Among all adults, the proportion who reported ever being told by a doctor that they had some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia. (N = 27,034)

Table 24: Cardiovascular Disease by EP Region
Michigan BRFSS, 2019-2021 Combined

	Ever Told Heart Attack ^a		Had Told Angina or Coronary Heart Disease ^b		Ever Told Stroke ^c		Ever Told Any Cardiovascular Disease ^d	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	4.7	(4.4-5.1)	4.6	(4.3-4.9)	3.6	(3.4-3.9)	9.7	(9.2-10.1)
EP Region								
Region 1	4.8	(4.0-5.8)	4.5	(3.8-5.3)	3.1	(2.5-3.8)	9.7	(8.6-11.0)
Region 2N	3.8	(3.3-4.5)	5.2	(4.5-5.9)	3.1	(2.6-3.6)	8.9	(8.1-9.9)
Region 2S	4.9	(4.2-5.7)	4.4	(3.8-5.2)	4.3	(3.7-5.1)	10.2	(9.2-11.3)
Region 3	6.4	(5.3-7.7)	4.9	(4.1-5.8)	4.9	(4.0-5.9)	11.7	(10.4-13.3)
Region 5	5.7	(4.7-7.0)	4.9	(4.0-6.0)	3.6	(2.9-4.6)	10.6	(9.2-12.1)
Region 6	3.6	(3.0-4.4)	3.6	(3.0-4.4)	2.8	(2.2-3.4)	7.4	(6.5-8.4)
Region 7	6.1	(4.8-7.8)	4.9	(3.7-6.5)	4.4	(3.3-6.0)	12.2	(10.3-14.4)
Region 8	6.1	(4.5-8.3)	7.6	(5.3-10.7)	3.4	(2.3-5.0)	12.7	(9.9-16.0)

Among all adults, the proportion who had ever been told by a doctor that...

^a they had a heart attack or myocardial infarction (N = 27,058);

^b they had angina or coronary heart disease (N = 26,975);

^c they had a stroke (N = 27,132);

^d they had a heart attack, coronary heart disease or a stroke (N = 26,935).

Table 25: Cancer by EP Region
Michigan BRFSS, 2019-2021 Combined

	Ever Told Skin Cancer ^a		Ever Told Any Other Types of Cancer ^b		Ever Told Cancer ^c	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	6.4	(6.1-6.7)	8.0	(7.7-8.4)	13.0	(12.5-13.5)
EP Region						
Region 1	7.0	(6.1-7.9)	8.8	(7.8-10.0)	14.1	(12.9-15.5)
Region 2N	7.0	(6.3-7.8)	8.0	(7.2-8.9)	13.5	(12.5-14.6)
Region 2S	5.0	(4.4-5.6)	7.6	(6.8-8.5)	11.4	(10.5-12.5)
Region 3	6.5	(5.6-7.6)	8.9	(7.8-10.2)	13.9	(12.5-15.5)
Region 5	7.7	(6.6-8.9)	8.4	(7.2-9.9)	14.2	(12.6-15.9)
Region 6	6.8	(6.1-7.6)	7.4	(6.6-8.4)	12.7	(11.6-13.9)
Region 7	7.1	(5.8-8.8)	8.5	(6.9-10.5)	14.5	(12.4-16.8)
Region 8	7.1	(5.4-9.2)	10.4	(7.9-13.5)	15.4	(12.6-18.7)

^a Among all adults, the proportion who reported ever being told by a doctor that they had skin cancer (N = 27,132).

^b Among all adults, the proportion who reported ever being told by a doctor that they had a form of cancer other than skin cancer (N = 27,131).

^c Among all adults, the proportion who reported ever being told by a doctor that they had skin cancer or any other type of cancer (N = 27,091).

Table 26: Diabetes by EP Region Michigan BRFSS, 2019-2021 Combined		
	Ever Told Diabetes^a	
	%	95% Confidence Interval
Michigan Total	11.4	(10.9-11.9)
EP Region		
Region 1	10.4	(9.3-11.6)
Region 2N	10.4	(9.4-11.5)
Region 2S	12.4	(11.3-13.5)
Region 3	14.2	(12.7-15.8)
Region 5	11.8	(10.4-13.4)
Region 6	11.1	(10.0-12.4)
Region 7	11.1	(9.2-13.3)
Region 8	12.4	(9.7-15.6)
^a Among all adults, the proportion who reported that they were ever told by a doctor that they had diabetes. Adults told they have prediabetes and women who had diabetes only during pregnancy were classified as not having been diagnosed. (N = 27,150)		

Table 27: Kidney Disease by EP Region Michigan BRFSS, 2019-2021 Combined		
	Ever Told Kidney Disease^a	
	%	95% Confidence Interval
Michigan Total	3.4	(3.1-3.6)
EP Region		
Region 1	3.2	(2.6-3.8)
Region 2N	3.0	(2.5-3.6)
Region 2S	3.6	(3.0-4.3)
Region 3	3.8	(3.1-4.6)
Region 5	4.2	(3.4-5.2)
Region 6	2.9	(2.4-3.6)
Region 7	3.7	(2.7-5.2)
Region 8	3.1	(1.9-5.0)

^a Among all adults, the proportion who reported ever being told by a doctor that they had kidney disease. (N = 27,117)

Table 28: Depression by EP Region Michigan BRFSS, 2019-2021 Combined		
	Ever Told Depression^a	
	%	95% Confidence Interval
Michigan Total	21.2	(20.5-21.9)
EP Region		
Region 1	22.9	(21.1-24.8)
Region 2N	18.1	(16.7-19.5)
Region 2S	20.7	(19.3-22.1)
Region 3	24.5	(22.4-26.8)
Region 5	23.2	(20.9-25.6)
Region 6	23.0	(21.2-25.0)
Region 7	19.4	(16.8-22.2)
Region 8	20.9	(17.6-24.6)

^a Among all adults, the proportion who reported ever being told by a doctor that they had a depressive disorder including depression, major depression, dysthymia, or minor depression. (N = 27,055)