

**Estimates for Selected Risk Factors and Health Indicators
by Emergency Preparedness Region**

State of Michigan

**Michigan Behavioral Risk Factor Surveillance System
(Michigan BRFSS)**

2020-2022

Lifecourse Epidemiology and Genomics Division
Bureau of Epidemiology and Population Health
Michigan Department of Health and Human Services

Table of Contents

Introduction	3
Map of Michigan Emergency Preparedness Regions	4
Health Status Tables	
Table 1: General Health Status	5
Table 2: Health Status on at Least 14 Days in the Past Month	6
Table 3: Mean Number of Days of Poor Physical Health, Poor Mental Health, and Activity Limitations.....	7
Table 4: Disability	8
Table 5: Weight Status	9
Table 6: Health Care Coverage Among Adults 18-64 Years Old	10
Table 7: Health Care Access	11
Risk Behaviors Tables	
Table 8: No Leisure Time Physical Activity	12
Table 9: Cigarette Smoking	13
Table 10: Current Smokers Who Attempted to Quit.....	14
Table 11: Hookah Use.....	15
Table 12: Alcohol Consumption.....	16
Table 13: Mean Number of Binge Drinking Occassions During the Past Month Among Binge Drinkers	17
Clinical Preventative Practices	
Table 14: No Routine Checkup in Past Year.....	18
Table 15: Breast Cancer Screening Among Women 40 Years and Older.....	19
Table 16: Cervical Cancer Screening Among Women 21-65 Years Old	20
Table 17: Colorectal Cancer Screening Among Adults 50 Years and Older	21
Table 18: Oral Health	22
Table 19: Immunizations Among Adults Aged 65 Years and Older	23
Table 20: HIV Testing Among Adults Aged 18 - 64 Years	24
Chronic Conditions	
Table 21: Asthma Among Adults	25
Table 22: Asthma Among Children.....	26
Table 23: Chronic Obstructive Pulmonary Disease (COPD).....	27
Table 24: Arthritis.....	28
Table 25: Cardiovascular Disease.....	29
Table 26: Cancer	30
Table 27: Diabetes	31
Table 28: Kidney Disease.....	32
Table 29: Depression	33

Introduction

The Michigan Behavioral Risk Factor Survey (MiBRFS) is an annual, statewide telephone survey of Michigan adults aged 18 years and older that is conducted through a collaborative effort among the Population Health Surveillance Branch (PHSB) of the Centers for Disease Control and Prevention (CDC), the ICF Macro Inc., and the Michigan Department of Health and Human Services. Michigan BRFSS data contribute to the national [Behavioral Risk Factor Surveillance System](#) that is managed by the PHSB at the CDC.

For this report, data from the 2020-2022 Michigan BRFSS were combined. The total sample sizes by year were: 7,269 in 2020, 9,425 in 2021, and 10,058 in 2022. Not all questions were included annually in the survey. The table subtitles indicate which years were used for that particular analysis.

For the 2020 through 2022 Michigan Behavioral Risk Factor Surveys, the sample of land line telephone numbers that were utilized for data collection was selected using a list-assisted, random-digit-dialed methodology with a disproportionate stratification based on phone bank density, and whether or not the phone numbers were directory listed. The sample of cell phone numbers used within each survey year was randomly selected from dedicated cellular telephone banks sorted on the basis of area code and exchange within the State of Michigan.

Iterative proportional fitting or raking was the weighting methodology used to make the 2020-2022 Michigan BRFSS data representative of the current Michigan adult population. Estimates based on this methodology were weighted to adjust for the probabilities of selection and a raking adjustment factor that adjusted for the distribution of Michigan adults by telephone source (land line vs. cell phone), detailed race/ethnicity, education level, marital status, age by gender, gender by race/ethnicity, age by race/ethnicity, and renter/owner status. No additional weighting factors were computed at the regional level, thus the weights used might not always provide accurate regional estimates, particularly when regional demographic distributions vary greatly from that of the state.

Michigan BRFSS data from 2020-2022 were analyzed by Michigan Emergency Preparedness (EP) region. A map identifying each of Michigan's eight emergency preparedness regions is included on the following page. Population-based prevalence estimates and asymmetric confidence intervals were calculated for indicators of health status, health risk behaviors, clinical preventive practices, and chronic conditions among the adult population in Michigan. All analyses were performed in SAS-Callable SUDAAN®, a statistical computing that is designed for complex sample surveys.

In an effort to provide the most reliable information possible, prevalence estimates are suppressed if they are based on a denominator of less than 50 respondents or have a relative standard error of greater than 30%.

Due to the BRFSS methodology changes that took place in 2011, the estimates provided within the following tables should not be compared to Michigan BRFSS estimates from years prior to 2011.

If you have any questions regarding any of the estimates within the following tables, please contact Yan Tian at TianY@michigan.gov.




Table 1: Health Status by EP Region Michigan BRFSS, 2020-2022 Combined		
	General Health, Fair or Poor^a	
	%	95% Confidence Interval
Michigan Total	16.3	(15.7-16.9)
EP Region		
Region 1	14.8	(13.4-16.4)
Region 2N	13.2	(12.1-14.5)
Region 2S	18.3	(17.0-19.7)
Region 3	19.2	(17.3-21.2)
Region 5	18.9	(16.9-21.1)
Region 6	15.3	(13.7-17.1)
Region 7	13.8	(11.4-16.7)
Region 8	18.1	(15.1-21.5)
^a Among all adults, the proportion who reported that their health, in general, was either fair or poor.		

Table 2: Health Status on at Least 14 Days in the Past Month by EP Region Michigan BRFSS, 2020-2022 Combined						
	Poor Physical Health^a		Poor Mental Health^b		Activity Limitation^c	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	11.9	(11.4-12.5)	16.0	(15.4-16.6)	9.7	(9.2-10.2)
EP Region						
Region 1	11.4	(10.1-12.9)	16.0	(14.3-17.7)	8.3	(7.2-9.6)
Region 2N	9.8	(8.9-10.9)	13.9	(12.7-15.3)	8.3	(7.4-9.3)
Region 2S	13.1	(11.9-14.4)	17.4	(16.1-18.8)	10.9	(9.8-12.0)
Region 3	14.4	(12.8-16.2)	18.3	(16.3-20.5)	11.4	(9.9-13.1)
Region 5	13.9	(12.1-15.9)	17.7	(15.5-20.1)	11.6	(9.8-13.6)
Region 6	11.2	(9.8-12.9)	14.9	(13.3-16.7)	8.9	(7.7-10.2)
Region 7	9.7	(7.7-12.0)	13.9	(11.3-16.9)	7.7	(6.0-9.9)
Region 8	14.2	(11.4-17.4)	13.1	(10.6-16.2)	9.4	(7.2-12.2)
<p>^a Among all adults, the proportion who reported 14 or more days of poor physical health, which includes physical illness and injury, during the past 30 days.</p> <p>^b Among all adults, the proportion who reported 14 or more days of poor mental health, which includes stress, depression, and problems with emotions, during the past 30 days.</p> <p>^c Among all adults, the proportion who reported 14 or more days in the past 30 days in which either poor physical health or poor mental health kept respondents from doing their usual activities, such as self-care, work, and recreation.</p>						

**Table 3: Mean Number of Days of Poor Physical Health, Poor Mental Health, and Activity Limitations by EP Region
Michigan BRFSS, 2020-2022 Combined**

	Poor Physical Health ^a		Poor Mental Health ^b		Activity Limitation ^c	
	Mean # of days	95% Confidence Interval	Mean # of days	95% Confidence Interval	Mean # of days	95% Confidence Interval
Michigan Total	3.8	(3.7-3.9)	5.1	(4.9-5.2)	3.1	(2.9-3.2)
EP Region						
Region 1	3.7	(3.3-4.0)	5.1	(4.6-5.5)	2.7	(2.4-3.0)
Region 2N	3.3	(3.0-3.6)	4.6	(4.3-4.9)	2.8	(2.6-3.1)
Region 2S	4.1	(3.8-4.4)	5.5	(5.2-5.8)	3.4	(3.2-3.7)
Region 3	4.3	(3.9-4.7)	5.6	(5.1-6.1)	3.6	(3.2-4.0)
Region 5	4.4	(3.9-4.9)	5.5	(4.9-6.0)	3.2	(2.8-3.6)
Region 6	3.7	(3.3-4.1)	4.7	(4.3-5.1)	2.8	(2.5-3.1)
Region 7	3.1	(2.6-3.6)	4.6	(3.9-5.3)	2.3	(1.9-2.8)
Region 8	4.4	(3.6-5.1)	4.5	(3.8-5.2)	3.0	(2.4-3.6)

^a Among all adults, the mean number of days during the past 30 days in which physical health, including illness and injury, was not good.

^b Among all adults, the mean number of days during the past 30 days in which mental health, including stress, depression, and problems with emotions, was not good.

^c Among all adults, the mean number of days during the past 30 days in which poor physical health or poor mental health kept respondent from doing their usual activities, such as self-care, work, and recreation.

Table 4: Disability by EP Region Michigan BRFSS, 2019 and 2020 Combined		
	Total Disability^a	
	%	95% Confidence Interval
Michigan Total	28.8	(28.0-29.5)
EP Region		
Region 1	27.6	(25.7-29.6)
Region 2N	24.2	(22.7-25.8)
Region 2S	30.8	(29.1-32.5)
Region 3	32.8	(30.5-35.3)
Region 5	32.8	(30.2-35.4)
Region 6	28.0	(26.0-30.1)
Region 7	30.9	(27.3-34.9)
Region 8	28.2	(24.4-32.2)
^a Among all adults, the proportion who reported having serious difficulty hearing, visual impairment, difficulty concentrating, remembering, or making decisions, difficulty walking or climbing stairs, difficulty dressing or bathing, or difficulty doing errands alone.		

Table 5: Weight Status by EP Region
Michigan BRFSS, 2020-2022 Combined

	Obese ^a		Overweight ^b		Healthy Weight ^c		Underweight ^d	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	34.7	(33.9-35.5)	33.9	(33.1-34.7)	29.3	(28.6-30.1)	2.1	(1.8-2.4)
EP Region								
Region 1	34.6	(32.4-36.7)	33.3	(31.2-35.5)	30.2	(28.1-32.3)	2.0	(1.4-2.9)
Region 2N	32.2	(30.5-34.0)	35.5	(33.7-37.3)	30.0	(28.3-31.8)	2.3	(1.7-3.2)
Region 2S	35.2	(33.5-37.0)	33.0	(31.3-34.7)	29.6	(28.0-31.3)	2.2	(1.6-2.9)
Region 3	41.1	(38.5-43.7)	33.7	(31.3-36.2)	22.4	(20.3-24.7)	2.8	(1.8-4.3)
Region 5	36.1	(33.6-38.8)	33.5	(31.0-36.0)	29.3	(26.7-32.0)	1.1	(0.7-1.8)
Region 6	35.2	(33.0-37.4)	34.6	(32.4-36.9)	28.8	(26.6-31.0)	1.4	(1.0-2.1)
Region 7	31.8	(28.3-35.6)	35.4	(31.7-39.2)	30.8	(27.3-34.6)	2.0	(1.1-3.5)
Region 8	34.2	(30.0-38.7)	33.6	(29.6-37.9)	30.9	(26.9-35.1)	-- ^e	-- ^e

Note: BMI, body mass index, is defined as weight (in kilograms) divided by height (in meters) squared [weight in kg/(height in meters)²]. Weight and height were self-reported. Pregnant women were excluded.

^a Among all adults, the proportion of respondents whose BMI was greater than or equal to 30.0.

^b Among all adults, the proportion of respondents whose BMI was greater than or equal to 25.0, but less than 30.0.

^c Among all adults, the proportion of respondents whose BMI was greater than or equal to 18.5, but less than 25.0.

^d Among all adults, the proportion of respondents whose BMI was less than 18.5.

^e Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 6: Health Care Coverage Among Adults 18-64 Years Old by EP Region Michigan BRFSS, 2020-2022 Combined		
	No Health Care Coverage^a (Among 18 - 64 year olds)	
	%	95% Confidence Interval
Michigan Total	6.8	(6.2-7.3)
EP Region		
Region 1	5.1	(4.0-6.5)
Region 2N	4.2	(3.3-5.2)
Region 2S	6.9	(5.8-8.2)
Region 3	7.5	(5.9-9.4)
Region 5	6.1	(4.7-7.9)
Region 6	7.8	(6.3-9.6)
Region 7	11.1	(6.9-17.3)
Region 8	6.0	(3.9-9.1)

^a Among adults aged 18-64 years, the proportion who reported having no health care coverage, including health insurance, prepaid plans such as HMOs, or government plans, such as Medicare or Indian Health Services.

Table 7: Health Care Access by EP Region Michigan BRFSS, 2020-2022 Combined				
	No Personal Health Care Provider^a		No Health Care Access During Past 12 Months Due to Cost^b	
	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	12.4	(11.8-13.0)	7.9	(7.5-8.4)
EP Region				
Region 1	12.5	(11.0-14.2)	7.4	(6.3-8.7)
Region 2N	9.4	(8.3-10.7)	6.2	(5.3-7.2)
Region 2S	12.9	(11.6-14.2)	8.1	(7.2-9.1)
Region 3	13.1	(11.2-15.1)	9.9	(8.4-11.6)
Region 5	12.8	(11.0-14.8)	8.2	(6.7-10.0)
Region 6	11.0	(9.5-12.7)	8.7	(7.4-10.2)
Region 7	12.4	(9.3-16.3)	5.7	(4.2-7.6)
Region 8	12.3	(9.6-15.6)	6.1	(4.4-8.3)
^a Among all adults, the proportion who reported that they did not have anyone that they thought of as their personal doctor or health care provider. ^b Among all adults, the proportion who reported that in the past 12 months, they could not see a doctor when they needed to due to the cost.				

Table 8: No Leisure Time Physical Activity by EP Region Michigan BRFSS, 2020-2022 Combined		
	No Leisure Time Physical Activity^a	
	%	95% Confidence Interval
Michigan Total	22.6	(22.0-23.3)
EP Region		
Region 1	21.9	(20.2-23.8)
Region 2N	20.1	(18.7-21.6)
Region 2S	24.2	(22.7-25.7)
Region 3	26.7	(24.6-28.9)
Region 5	23.4	(21.4-25.6)
Region 6	20.4	(18.7-22.2)
Region 7	22.4	(18.9-26.3)
Region 8	23.0	(19.9-26.6)

^a Among all adults, the proportion who reported not participating in any leisure time physical activities or exercises such as running, calisthenics, golf, gardening, or walking during the past month.

Table 9: Cigarette Smoking by EP Region
Michigan BRFSS, 2020-2022 Combined

	Current Smoking ^a		Former Smoking ^b		Never Smoked	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	16.9	(16.3-17.6)	26.5	(25.8-27.3)	56.6	(55.7-57.4)
EP Region						
Region 1	16.4	(14.8-18.2)	28.4	(26.5-30.4)	55.2	(53.0-57.4)
Region 2N	14.3	(12.9-15.7)	26.6	(25.0-28.2)	59.2	(57.3-61.0)
Region 2S	16.8	(15.4-18.2)	23.9	(22.5-25.4)	59.3	(57.5-61.0)
Region 3	20.3	(18.3-22.5)	29.8	(27.4-32.2)	49.9	(47.3-52.5)
Region 5	19.0	(16.7-21.4)	27.8	(25.6-30.1)	53.2	(50.6-55.9)
Region 6	17.1	(15.2-19.2)	26.2	(24.3-28.2)	56.7	(54.4-58.9)
Region 7	18.7	(15.6-22.2)	28.2	(24.9-31.8)	53.1	(49.0-57.1)
Region 8	18.5	(15.4-22.0)	30.4	(26.8-34.3)	51.1	(46.9-55.3)

^a Among all adults, the proportion who reported that they had ever smoked at least 100 cigarettes (5 packs) in their life and that they smoke cigarettes now, either every day or on some days.

^b Among all adults, the proportion who reported that they had ever smoked at least 100 cigarettes (5 packs) in their life, but they do not smoke cigarettes now.

Table 10: Current Smokers Who Attempted to Quit by EP Region Michigan BRFSS, 2020-2022 Combined		
	Tried to Quit Smoking One Day or Longer in Past Year^a	
	%	95% Confidence Interval
Michigan Total	55.2	(52.9-57.6)
EP Region		
Region 1	53.1	(46.8-59.2)
Region 2N	56.3	(50.6-61.8)
Region 2S	61.9	(56.9-66.6)
Region 3	53.2	(47.1-59.2)
Region 5	56.7	(49.2-63.9)
Region 6	50.9	(43.9-57.9)
Region 7	49.4	(39.5-59.5)
Region 8	51.3	(40.7-61.9)

^a Among current smokers, the proportion who reported that during the past 12 months, they had tried to quit smoking for one day or longer.

Table 11: Current Hookah Use by EP Region Michigan BRFSS, 2020-2022 Combined		
	Current Hookah Use^a	
	%	95% Confidence Interval
Michigan Total	2.0	(1.7-2.4)
EP Region		
Region 1	1.6	(1.0-2.6)
Region 2N	2.3	(1.6-3.3)
Region 2S	3.2	(2.4-4.3)
Region 3	1.7	(1.0-2.8)
Region 5	-- ^b	-- ^b
Region 6	1.2	(0.7-1.9)
Region 7	-- ^b	-- ^b
Region 8	-- ^b	-- ^b

^a Among all adults, the proportion who reported smoking tobacco using a hookah, narghile, or water pipe on one or more days during the previous thirty day.
^b Suppressed due to a denominator < 50 and/or a relative standard error > 30%.

Table 12: Alcohol Consumption by EP Region
Michigan BRFSS, 2020-2022 Combined

	Any Alcohol in Past Month ^a		Heavy Drinking ^b		Binge Drinking ^c	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	54.8	(54.0-55.6)	6.8	(6.4-7.2)	16.9	(16.2-17.5)
EP Region						
Region 1	54.2	(51.9-56.4)	6.3	(5.3-7.5)	17.2	(15.5-19.1)
Region 2N	59.9	(58.0-61.7)	6.4	(5.6-7.4)	16.4	(15.0-17.9)
Region 2S	52.3	(50.5-54.1)	6.1	(5.3-7.0)	16.1	(14.7-17.5)
Region 3	51.9	(49.3-54.4)	6.6	(5.5-8.0)	17.8	(15.7-20.1)
Region 5	52.7	(50.1-55.4)	7.1	(5.8-8.7)	14.7	(12.9-16.8)
Region 6	55.5	(53.1-57.8)	7.5	(6.2-9.1)	17.3	(15.5-19.2)
Region 7	56.2	(52.0-60.3)	8.3	(6.4-10.6)	16.1	(13.4-19.2)
Region 8	55.2	(50.9-59.4)	8.8	(6.6-11.6)	22.3	(18.7-26.2)

^a Among all adults, the proportion who reported consuming at least one drink of any alcoholic beverage in the previous month.
^b Among all adults, the proportion who reported consuming an average of more than two alcoholic drinks per day for men or more than one per day for women in the previous month.
^c Among all adults, the proportion who reported consuming five or more drinks per occasion (for men) or 4 or more drinks per occasion (for women) at least once in the previous month.

Table 13: Mean Number of Binge Drinking Occasions during the Past Month among Binge Drinkers by EP Region Michigan BRFSS, 2020-2022 Combined		
	Number of Binge Drinking Occasions^a	
	Mean	95% Confidence Interval
Michigan Total	5.2	(4.8-5.5)
EP Region		
Region 1	4.1	(3.5-4.7)
Region 2N	4.6	(3.8-5.3)
Region 2S	5.0	(4.3-5.7)
Region 3	5.1	(4.1-6.1)
Region 5	5.3	(4.4-6.3)
Region 6	5.9	(4.8-7.1)
Region 7	7.0	(5.1-9.0)
Region 8	5.3	(3.9-6.7)

^a Among adults who binge drank in the past month, the mean number of times during which five or more alcoholic beverages (for men) or four or more (for women) were consumed during the past month.

Table 14: No Routine Checkup in Past Year by EP Region Michigan BRFSS, 2020-2022 Combined		
	Had No Routine Checkup in Past Year^a	
	%	95% Confidence Interval
Michigan Total	22.3	(21.6-23.0)
EP Region		
Region 1	22.7	(20.9-24.7)
Region 2N	21.4	(19.9-23.0)
Region 2S	21.9	(20.5-23.4)
Region 3	21.4	(19.4-23.6)
Region 5	23.4	(21.2-25.8)
Region 6	23.2	(21.3-25.3)
Region 7	19.0	(15.5-23.0)
Region 8	21.4	(18.0-25.2)

^a Among all adults, the proportion who reported that they did not have a routine checkup in the past year.

Table 15: Breast Cancer Screening Among Women Aged 40 Years and Older by EP Region Michigan BRFSS, 2020 and 2022 Combined		
	Had Mammogram in Past Two Years^a	
	%	95% Confidence Interval
Michigan Total	72.7	(71.3-74.1)
EP Region		
Region 1	76.3	(72.7-79.6)
Region 2N	74.1	(70.8-77.0)
Region 2S	69.5	(66.1-72.7)
Region 3	74.9	(70.5-78.9)
Region 5	70.5	(66.2-74.6)
Region 6	72.8	(68.8-76.4)
Region 7	75.6	(68.8-81.2)
Region 8	74.5	(67.7-80.2)

^a Among women aged 40 years and older, the proportion who reported having a mammogram in the past two years.

Table 16: Cervical Cancer Screening Among Women Aged 21-65 Years Old by EP Region Michigan BRFSS, 2022		
	Had Appropriately Timed Cervical Cancer Screening Test^a	
	%	95% Confidence Interval
Michigan Total	54.6	(51.9-57.4)
EP Region		
Region 1	57.1	(49.1-64.8)
Region 2N	60.0	(53.8-66.0)
Region 2S	51.5	(45.9-57.1)
Region 3	53.1	(44.2-61.8)
Region 5	51.7	(42.0-61.2)
Region 6	56.2	(48.4-63.7)
Region 7	67.7	(54.4-78.7)
Region 8	58.4	(42.0-73.1)

^a Among women aged 21-65 years old, the proportion who reported having a cervical cancer screening test within the previous three years.

Table 17: Colorectal Cancer Screening Among Adults Aged 50 Years and Older by EP Region Michigan BRFSS, 2020 and 2022 Combined		
	Appropriate Colorectal Cancer Screening^a	
	%	95% Confidence Interval
Michigan Total	75.4	(74.3-76.5)
EP Region		
Region 1	79.1	(76.3-81.6)
Region 2N	75.9	(73.3-78.4)
Region 2S	74.3	(71.7-76.7)
Region 3	76.8	(73.3-80.0)
Region 5	76.4	(73.1-79.4)
Region 6	74.8	(71.5-77.9)
Region 7	73.9	(67.9-79.1)
Region 8	78.4	(73.2-82.8)
^a Among adults aged 50 years and older, the proportion who reported having either a fecal occult blood test within the past year, a sigmoidoscopy within the past five years, or a colonoscopy within the past ten years.		

Table 18: Oral Health by EP Region Michigan BRFSS, 2020 and 2022 Combined				
	No Dental Visit in Past Year^a		6+ Teeth Missing^b	
	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	30.7	(29.8-31.7)	14.2	(13.5-14.9)
EP Region				
Region 1	28.3	(26.0-30.8)	13.7	(12.0-15.6)
Region 2N	25.9	(23.9-28.0)	10.5	(9.3-11.9)
Region 2S	32.3	(30.3-34.4)	14.3	(12.9-15.9)
Region 3	34.6	(31.5-37.8)	17.4	(15.3-19.7)
Region 5	31.6	(28.5-34.8)	17.1	(14.8-19.7)
Region 6	30.2	(27.7-33.0)	14.1	(12.1-16.4)
Region 7	29.9	(24.9-35.3)	16.2	(12.0-21.5)
Region 8	33.4	(28.3-39.0)	24.1	(19.5-29.4)

^a Among all adults, the proportion who reported that they had not visited a dentist or dental clinic for any reason within the previous year.

^b Among all adults, the proportion who reported that they were missing 6+ teeth due to tooth decay or gum disease. This excludes teeth lost for other reasons, such as injury or orthodontics.

**Table 19: Immunizations among Adults Aged 65 Years and Older
by EP Region**

Michigan BRFSS, 2020-2022 Combined

	Had Flu Vaccine in Past Year ^a		Ever Had Pneumonia Vaccine ^b	
	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	72.0	(70.8-73.2)	72.8	(71.5-74.0)
EP Region				
Region 1	72.4	(69.3-75.3)	75.8	(72.8-78.6)
Region 2N	74.6	(71.9-77.2)	74.1	(71.4-76.7)
Region 2S	71.7	(68.7-74.6)	69.6	(66.4-72.6)
Region 3	64.3	(60.2-68.3)	68.2	(64.0-72.1)
Region 5	75.7	(72.2-78.9)	75.1	(71.4-78.4)
Region 6	74.5	(71.2-77.6)	75.0	(71.7-78.1)
Region 7	74.9	(68.6-80.4)	72.4	(66.0-77.9)
Region 8	72.3	(65.8-78.1)	76.9	(70.3-82.5)

^a Among adults aged 65 years and older, the proportion who reported that they had a flu vaccine, either by an injection in the arm or sprayed in the nose during the past 12 months.

^b Among adults aged 65 years and older, the proportion who reported that they ever had a pneumococcal vaccine.

Table 20: HIV Testing among Adults Aged 18 - 64 Years by EP Region Michigan BRFSS, 2020-2022 Combined		
	Ever Had an HIV Test^a	
	%	95% Confidence Interval
Michigan Total	43.5	(42.5-44.6)
EP Region		
Region 1	43.2	(40.3-46.0)
Region 2N	43.3	(40.9-45.6)
Region 2S	51.1	(48.9-53.3)
Region 3	39.1	(35.8-42.4)
Region 5	47.0	(43.5-50.6)
Region 6	39.6	(36.6-42.5)
Region 7	40.2	(35.0-45.6)
Region 8	34.9	(29.9-40.3)

^a Among adults aged 18 - 64 years, the proportion who reported that they ever had been tested for HIV, apart from tests that were part of a blood donation. Don't know was considered a valid response to this question.

Table 21: Asthma among Adults by EP Region Michigan BRFSS, 2020-2022 Combined				
	Lifetime Asthma Prevalence^a		Current Asthma Prevalence^b	
	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	16.2	(15.6-16.8)	11.5	(11.0-12.0)
EP Region				
Region 1	17.1	(15.5-18.9)	12.7	(11.3-14.3)
Region 2N	16.3	(15.0-17.7)	10.8	(9.7-12.0)
Region 2S	18.1	(16.8-19.5)	13.1	(12.0-14.3)
Region 3	15.2	(13.4-17.1)	11.1	(9.6-12.8)
Region 5	16.5	(14.6-18.7)	12.5	(10.7-14.5)
Region 6	14.1	(12.6-15.8)	9.6	(8.4-10.9)
Region 7	15.5	(12.2-19.5)	11.7	(8.7-15.6)
Region 8	16.6	(13.8-19.9)	11.6	(9.4-14.4)
^a Among all adults, the proportion who reported that they were ever told by a doctor, nurse, or other health care professional that they had asthma. ^b Among all adults, the proportion who reported that they still had asthma.				

Table 22: Asthma among Children by EP Region Michigan BRFSS, 2020-2022 Combined				
	Lifetime Asthma Prevalence^a		Current Asthma Prevalence^b	
	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	10.9	(9.7-12.1)	7.4	(6.4-8.5)
EP Region				
Region 1	10.9	(8.4-14.0)	7.1	(5.2-9.7)
Region 2N	12.0	(9.2-15.5)	7.7	(5.4-10.9)
Region 2S	12.0	(9.7-14.8)	7.9	(6.2-10.2)
Region 3	11.3	(8.1-15.7)	7.8	(4.9-12.1)
Region 5	10.5	(7.3-15.0)	7.7	(5.1-11.6)
Region 6	10.1	(7.4-13.7)	6.6	(4.6-9.5)
Region 7	4.8	(2.7-8.5)	-- ^c	-- ^c
Region 8	-- ^c	-- ^c	-- ^c	-- ^c
^a Estimated proportion of Michigan children aged 0-17 years who were ever told by a doctor, nurse or other health professional that they had asthma, using proxy information from the adult respondent. ^b Estimated proportion of Michigan children aged 0-17 years who still have asthma, using proxy information from the adult respondent. ^c Suppressed due to a denominator < 50 and/or a relative standard error > 30%.				

Table 23: Chronic Obstructive Pulmonary Disease (COPD) by EP Region Michigan BRFSS, 2020-2022 Combined		
	Ever Told COPD, Emphysema or Chronic Bronchitis^a	
	%	95% Confidence Interval
Michigan Total	8.3	(7.9-8.8)
EP Region		
Region 1	9.1	(7.9-10.4)
Region 2N	7.4	(6.5-8.3)
Region 2S	8.8	(7.9-9.8)
Region 3	11.1	(9.7-12.7)
Region 5	8.7	(7.3-10.3)
Region 6	7.1	(6.1-8.2)
Region 7	7.4	(5.7-9.5)
Region 8	9.4	(7.2-12.2)

^a Among all adults, the proportion who reported ever being told by a doctor that they had chronic obstructive pulmonary disease (COPD), emphysema or chronic bronchitis.

Table 24: Arthritis by EP Region Michigan BRFSS, 2020-2022 Combined		
	Ever Told Arthritis^a	
	%	95% Confidence Interval
Michigan Total	30.0	(29.3-30.7)
EP Region		
Region 1	31.4	(29.6-33.4)
Region 2N	28.2	(26.7-29.7)
Region 2S	29.6	(28.1-31.1)
Region 3	36.0	(33.8-38.3)
Region 5	33.0	(30.7-35.4)
Region 6	28.9	(27.1-30.8)
Region 7	32.4	(29.0-35.9)
Region 8	35.7	(32.0-39.7)

^a Among all adults, the proportion who reported ever being told by a doctor that they had some form of arthritis, rheumatoid arthritis, gout, lupus, or fibromyalgia.

Table 25: Cardiovascular Disease by EP Region
Michigan BRFSS, 2020-2022 Combined

	Ever Told Heart Attack ^a		Had Told Angina or Coronary Heart Disease ^b		Ever Told Stroke ^c		Ever Told Any Cardiovascular Disease ^d	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	4.8	(4.5-5.2)	4.6	(4.3-4.9)	3.6	(3.3-3.8)	9.8	(9.4-10.3)
EP Region								
Region 1	5.2	(4.4-6.3)	4.9	(4.2-5.8)	3.0	(2.4-3.7)	10.3	(9.1-11.6)
Region 2N	4.0	(3.3-4.7)	4.9	(4.3-5.6)	2.9	(2.4-3.4)	9.1	(8.2-10.1)
Region 2S	5.1	(4.4-5.9)	4.5	(3.9-5.3)	4.3	(3.6-5.0)	10.2	(9.2-11.3)
Region 3	6.0	(4.9-7.2)	4.8	(4.0-5.7)	4.6	(3.7-5.6)	11.7	(10.3-13.2)
Region 5	6.0	(4.8-7.4)	5.2	(4.1-6.5)	4.0	(3.2-5.0)	11.1	(9.6-12.8)
Region 6	3.8	(3.2-4.7)	4.1	(3.4-4.9)	3.0	(2.4-3.7)	8.0	(7.1-9.1)
Region 7	5.5	(4.1-7.3)	4.6	(3.4-6.2)	3.8	(2.8-5.2)	10.8	(8.9-13.0)
Region 8	6.1	(4.6-8.1)	6.6	(4.6-9.6)	3.2	(2.2-4.7)	12.3	(9.7-15.5)

Among all adults, the proportion who had ever been told by a doctor that...

^a they had a heart attack or myocardial infarction;

^b they had angina or coronary heart disease;

^c they had a stroke;

^d they had a heart attack, coronary heart disease or a stroke.

Table 26: Cancer by EP Region
Michigan BRFSS, 2020-2022 Combined

	Ever Told Skin Cancer ^a		Ever Told Any Other Types of Cancer ^b		Ever Told Cancer ^c	
	%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
Michigan Total	5.9	(5.6-6.2)	8.1	(7.8-8.5)	12.4	(12.0-12.9)
EP Region						
Region 1	6.5	(5.7-7.4)	9.7	(8.6-10.9)	14.4	(13.2-15.8)
Region 2N	6.0	(5.3-6.7)	8.2	(7.4-9.1)	12.6	(11.6-13.7)
Region 2S	4.6	(4.1-5.2)	7.1	(6.4-7.9)	10.5	(9.6-11.4)
Region 3	6.2	(5.2-7.2)	8.8	(7.6-10.1)	13.2	(11.8-14.7)
Region 5	7.5	(6.5-8.7)	9.5	(8.1-11.1)	14.8	(13.1-16.6)
Region 6	6.6	(5.9-7.5)	7.8	(6.9-8.8)	12.6	(11.5-13.8)
Region 7	6.9	(5.5-8.5)	9.0	(7.3-11.0)	14.3	(12.2-16.7)
Region 8	6.7	(5.2-8.4)	10.3	(7.9-13.2)	14.7	(12.1-17.7)

^a Among all adults, the proportion who reported ever being told by a doctor that they had skin cancer.

^b Among all adults, the proportion who reported ever being told by a doctor that they had a form of cancer other than skin cancer.

^c Among all adults, the proportion who reported ever being told by a doctor that they had skin cancer or any other type of cancer.

Table 27: Diabetes by EP Region Michigan BRFSS, 2020-2022 Combined		
	Ever Told Diabetes^a	
	%	95% Confidence Interval
Michigan Total	11.6	(11.1-12.1)
EP Region		
Region 1	10.7	(9.6-12.0)
Region 2N	10.4	(9.4-11.5)
Region 2S	13.4	(12.3-14.6)
Region 3	14.8	(13.2-16.5)
Region 5	11.4	(10.0-12.9)
Region 6	11.2	(10.0-12.6)
Region 7	9.8	(8.1-11.9)
Region 8	11.8	(9.3-14.9)

^a Among all adults, the proportion who reported that they were ever told by a doctor that they had diabetes. Adults told they have prediabetes and women who had diabetes only during pregnancy were classified as not having been diagnosed.

Table 28: Kidney Disease by EP Region Michigan BRFSS, 2020-2022 Combined		
	Ever Told Kidney Disease^a	
	%	95% Confidence Interval
Michigan Total	3.6	(3.3-3.9)
EP Region		
Region 1	3.3	(2.7-4.1)
Region 2N	3.0	(2.5-3.6)
Region 2S	3.7	(3.1-4.3)
Region 3	4.0	(3.3-4.9)
Region 5	4.1	(3.3-5.1)
Region 6	3.8	(3.1-4.7)
Region 7	3.6	(2.5-5.2)
Region 8	3.6	(2.3-5.7)

^a Among all adults, the proportion who reported ever being told by a doctor that they had kidney disease.

Table 29: Depression by EP Region Michigan BRFSS, 2020-2022 Combined		
	Ever Told Depression^a	
	%	95% Confidence Interval
Michigan Total	21.7	(21.1-22.4)
EP Region		
Region 1	22.6	(20.8-24.5)
Region 2N	18.1	(16.7-19.5)
Region 2S	21.2	(19.8-22.6)
Region 3	25.4	(23.2-27.7)
Region 5	24.5	(22.2-27.0)
Region 6	23.1	(21.2-25.1)
Region 7	18.5	(15.8-21.4)
Region 8	22.0	(18.7-25.5)

^a Among all adults, the proportion who reported ever being told by a doctor that they had a depressive disorder including depression, major depression, dysthymia, or minor depression.