



NOW IT'S YOUR TURN TO

**CRUSH
HEP C**

**WHAT YOU NEED TO KNOW
TO GET CURED*
OF CHRONIC HEPATITIS C**

*Cure means no hep C virus found in the blood 3 months after treatment ends. Individual results may vary.

CONGRATS! YOU'RE ON YOUR WAY TO CURE

Take MAVYRET for only

8
WEEKS

For those new to treatment



When will you know that you're CURED?

Your doctor will do a blood test 12 weeks after you finish taking MAVYRET, and that's when you'll learn if you've been CURED

#1

**PRESCRIBED MEDICINE
FOR CHRONIC HEPATITIS C***

*Based on data from IQVIA, week ending 1/12/2018 to 8/11/2023.

USE

MAVYRET is a prescription medicine used to treat adults and children 3 years of age and older with chronic (lasting a long time) hepatitis C virus (hep C) genotypes 1, 2, 3, 4, 5 or 6 infection without cirrhosis or with compensated cirrhosis.

SAFETY CONSIDERATIONS

Hepatitis B virus (hep B) reactivation: Before starting treatment with MAVYRET, your doctor will do blood tests to check for hep B infection. If you have ever had hep B infection, hep B could become active again during or after treatment for hep C with MAVYRET. Hep B that becomes active again (called reactivation) may cause serious liver problems, including liver failure and death. Your doctor will monitor you if you are at risk for hep B reactivation during treatment and after you stop taking MAVYRET.

AFTER ONLY 8 WEEKS, YOU TOO CAN KNOW HOW IT FEELS TO BE CURED



The day I found out I was cured I was on top of the world.

**BEING CURED
JUST FEELS AMAZING.**

First thing I did was
run out of the house—
'I'M CURED!'



**FINDING OUT
I WAS CURED**

was the spark
of hope I needed.



These are all real MAVYRET patients.

WHAT WILL YOUR STORY OF CURE BE?

When you're **CURED** and ready to share your story, give us a call at **1.800.324.5074** or email us at info@Speak-MA.net

Please see additional Important Safety Information on pages 10 and 11.

Please see full Prescribing Information, including the Patient Information.

MAVYRET
glecaprevir/pibrentasvir
100 mg/40 mg tablets

TAKING MAVYRET IS SIMPLE



Once a day – 3 pills at one time – with food*

- MAVYRET can fit into your daily life because you take it only once a day



What to remember while taking MAVYRET

- Always take MAVYRET exactly as your doctor tells you
- Don't change, miss, or skip doses
- Do not stop taking MAVYRET without talking to your doctor first
- Always take MAVYRET with food

If you take too much MAVYRET, call your doctor or go to the nearest hospital emergency room right away.

What to do if you miss a dose

If it's less than 18 hours:

Take the missed dose with food as soon as you can. Then take your next dose at your usual time.



If it's more than 18 hours:

Do not take your missed dose. Take your next dose, as usual, with food.



*For adults, adolescents 12 years and older, or children weighing at least 99 pounds (45 kg).



**Taking MAVYRET WAS
EASY FOR ME.**

**I didn't have to worry
about it—it was great.**



Dianna, CURED with MAVYRET

SAFETY CONSIDERATIONS (cont'd)

- Do not take MAVYRET if you have certain liver problems or if you are taking the medicines atazanavir or rifampin.

Please see additional Important Safety Information, including BOXED WARNING on Hep B Reactivation, on pages 10 and 11.

Please see full Prescribing Information, including the Patient Information.

MAVYRET 
glecaprevir/pibrentasvir
100 mg/40 mg tablets

ANSWERS TO COMMON QUESTIONS



What are the common side effects of MAVYRET?

The most common side effects are headaches and tiredness. These are not all the possible side effects of MAVYRET. Tell your doctor if you have any side effect that bothers you or that does not go away.



What's important to know about the side effects of MAVYRET?

- Most side effects were mild
- Only 3 out of ~2,300 (0.1%) people in clinical trials stopped taking MAVYRET due to side effects

You may report side effects to the FDA at [1-800-FDA-1088](tel:1-800-FDA-1088) or by visiting www.fda.gov/medwatch.

SAFETY CONSIDERATIONS (cont'd)

In people who had or have advanced liver problems before starting treatment with MAVYRET, there is a rare risk of worsening liver problems, liver failure, and death. Your doctor will check you for signs and symptoms of worsening liver problems during treatment with MAVYRET. Tell your doctor right away if you have any of the following: nausea; tiredness; yellowing of your skin or white part of your eyes; bleeding or bruising more easily than normal; confusion; dark, black, or bloody stool; loss of appetite; diarrhea; dark or brown (tea-colored) urine; swelling or pain on the upper right side of your stomach area (abdomen); sleepiness; vomiting of blood; or lightheadedness.

Please see additional Important Safety Information, including BOXED WARNING on Hep B Reactivation, on pages 10 and 11.



What about pregnancy?

It's important to tell your doctor if you become pregnant during treatment with MAVYRET.

Since it's not known if MAVYRET passes into breast milk, your doctor can tell you the best way to feed your baby when you are on treatment.



Can I take other medicines while I'm taking MAVYRET?

You can take MAVYRET while taking many other medicines, including:

- Medication-assisted treatment for opioid-use disorder*
- Many types of birth control—but make sure to talk to your doctor

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

*If you are taking naltrexone, talk to your doctor to learn if you can still take MAVYRET.

Please see full Prescribing Information, including the Patient Information.

MAVYRET
glecaprevir/pibrentasvir
100 mg/40 mg tablets

HELPFUL INFORMATION ON INSURANCE

Commercial insurance?

Claim your MAVYRET Savings Card today—
you may pay as little as \$5 a month*

SCAN TO GET THE
SAVINGS CARD TODAY



Medicaid?

Pay \$20 or less, depending on your state plan

Uninsured or having difficulty paying for your medication?



myAbbVie Assist provides AbbVie
medicines to qualifying patients.
[Click here](#) or call **1-800-222-6885**
to learn more.

*Eligibility: Available to patients with commercial insurance coverage for MAVYRET who meet eligibility criteria. This co-pay assistance program is not available to patients receiving prescription reimbursement under any federal, state, or government-funded insurance programs (for example, Medicare [including Part D], Medicare Advantage, Medigap, Medicaid, TRICARE, Department of Defense, or Veterans Affairs programs) or where prohibited by law. Offer subject to change or termination without notice. Restrictions, including monthly maximums, may apply. This is not health insurance. For full Terms and Conditions, visit <https://www.mavyret.com/cost> or call **877-628-9738** for additional information. To learn about AbbVie's privacy practices and your privacy choices, visit <https://privacy.abbvie>

THE SUPPORT YOU NEED, WHEN YOU NEED IT



Once you have your prescription, the MAVYRET Patient Support program will be there to support you every step of the way

From the start, you'll be connected with your own Nurse Ambassador,* to provide resources and support throughout your treatment, including:

- answering your questions about MAVYRET
- identifying possible ways to save on MAVYRET
- making sense of your insurance coverage



To get started with MAVYRET Patient Support, call **1.877.MAVYRET** or scan to visit mavyret.com/patient-support

*Ambassadors are provided by AbbVie and do not provide medical advice or work under the direction of the prescribing health care professional (HCP). They are trained to direct patients to speak with their HCP about any treatment-related questions, including further referrals.

Please see additional Important Safety Information, including **BOXED WARNING** on Hep B Reactivation, on pages 10 and 11.

Please see full Prescribing Information, including the Patient Information.

MAVYRET 
glecaprevir/pibrentasvir
100 mg/40 mg tablets

IMPORTANT SAFETY INFORMATION AND USE

What is the most important information I should know about MAVYRET?

Hepatitis B virus (hep B) reactivation: Before starting treatment with MAVYRET, your doctor will do blood tests to check for hep B infection. If you have ever had hep B infection, hep B could become active again during or after treatment for hep C with MAVYRET. Hep B that becomes active again (called reactivation) may cause serious liver problems, including liver failure and death. Your doctor will monitor you if you are at risk for hep B reactivation during treatment and after you stop taking MAVYRET.

Do not take MAVYRET if you:

- Have certain liver problems
- Are taking the medicines atazanavir or rifampin

What should I tell my doctor before taking MAVYRET?

- If you have had hep B infection, have liver problems other than hep C infection, have HIV-1 infection, have had a liver or a kidney transplant, and all other medical conditions.
- If you are pregnant or plan to become pregnant, or if you are breastfeeding or plan to breastfeed. It is not known if MAVYRET will harm your unborn baby or pass into your breast milk. Talk to your doctor about the best way to feed your baby if you take MAVYRET.
- **About all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. MAVYRET and other medicines may affect each other. This can cause you to have too much or not enough MAVYRET or other medicines in your body. This may affect the way MAVYRET or your other medicines work or may cause side effects.
- **Do not start taking a new medicine without telling your doctor.** Your doctor can tell you if it is safe to take MAVYRET with other medicines.

What are the possible side effects of MAVYRET?

- **In people who had or have advanced liver problems before starting treatment with MAVYRET, there is a rare risk of worsening liver problems, liver failure, and death.** Your doctor will check you for signs and symptoms of worsening liver problems during treatment with MAVYRET. Tell your doctor right away if you have any of the following: nausea; tiredness; yellowing of your skin or white part of your eyes; bleeding or bruising more easily than normal; confusion; dark, black, or bloody stool; loss of appetite; diarrhea; dark or brown (tea-colored) urine; swelling or pain on the upper right side of your stomach area (abdomen); sleepiness; vomiting of blood; or lightheadedness.

- The most common side effects of MAVYRET are headache and tiredness.

These are not all the possible side effects of MAVYRET. Call your doctor for medical advice about side effects.

This is the most important information to know about MAVYRET. For more information, talk to your doctor or healthcare provider.

MAVYRET oral pellets are dispensed in unit-dose packets. Each packet contains 50 mg glecaprevir/20 mg pibrentasvir.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call [1-800-FDA-1088](tel:1-800-FDA-1088).

USE

MAVYRET is a prescription medicine used to treat adults and children 3 years of age and older with chronic (lasting a long time) hepatitis C virus (hep C):

- Genotypes (GT) 1, 2, 3, 4, 5 or 6 infection without cirrhosis or with compensated cirrhosis
- Or GT 1 infection and have been previously treated with a regimen that contained a hep C NS5A inhibitor or an NS3/4A protease inhibitor, but not both.

Please see full Prescribing Information, including the Patient Information.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/myAbbVieAssist to learn more.

“**BEING CURED**
was a life changer
for me.”

**Danny, CURED
with MAVYRET**

MAVYRET
glecaprevir/pibrentasvir
100 mg/40 mg tablets

TREATMENT MILESTONES ON THE WAY TO YOUR CURE

Start treatment: ____/____/____

Finish treatment: ____/____/____

Officially **CURED***: ____/____/____
(after 12-week test results)

*Cure means no hep C virus found in the blood 3 months after treatment ends.

If you have any questions for your healthcare professional, write them down here for your next visit



If you have any questions while taking MAVYRET, talk to your doctor or call 1.877.MAVYRET

SAFETY CONSIDERATIONS (cont'd)

- Do not take MAVYRET if you have certain liver problems or if you are taking the medicines atazanavir or rifampin.

Please see additional Important Safety Information, including **BOXED WARNING** on Hep B Reactivation, on pages 10 and 11.

Please see full Prescribing Information, including the Patient Information.

abbvie

MAVYRET[®]
glecaprevir/pibrentasvir
100 mg/40 mg tablets